ACCEL CALF ADE+ TYPE B MEDICATED FEED

A calf supplement to be mixed with dry non-medicated milk replacer to make a complete medicated Type C milk replacer ration for calves and beef and non-lactating dairy cattle.

FEATURES

Provides each calf with 10 mg. of oxytetracycline and 10 mg. of neomycin sulfate per pound of calf body weight daily.

BENEFITS

For the treatment of bacterial enteritis (scours) caused by Escherichia coli and bacterial pneumonia (shipping fever complex) caused by Pasteurella multocida susceptible to oxytetracycline; also for the treatment and control of colibacillosis (bacterial enteritis, scours) caused by Escherichia coli susceptible to neomycin. Feed to provide each calf *10 mg of oxytetracycline and 10 mg of neomycin sulfate per pound of calf bodyweight daily. Feed continously for 7 to 14 days in place of normal milk replacer.

APPLICATION

- Mix the entire package (4.34 lbs.) of Accel Calf Ade+ powder into 50 pounds of dry non-medicated calf milk replacer
 powder to make a Type C complete, medicated milk replacer ration containing 1,000 mg Oxytetracycline and 1,000 mg
 Neomycin Sulfate per 1.25 pounds of medicated milk replacer powder.
- Mix and feed the resulting Type C feed as a complete ration according to the following chart:

Calf Weight (lbs.)	*Neomycin & Oxytetracycline (mg/day)	Type C Milk Replacer Powder (ounces)	Water (115-120°F) (quarts)	Feedings Per Day
80	800	8	2	2
100	1000	10	2	2
125	1250	12.5	2.5	2
150	1500	15	3	2

If symptoms persist after using for 2 or 3 days, consult a veterinarian. Treatment should continue 24 to 48 hours beyond remission of disease symptoms.

FOR USE IN DRY FEEDS ONLY. NOT FOR USE IN LIQUID FEED SUPPLEMENTS. FOR USE IN ANIMAL FEED ONLY. NOT FOR HUMAN USE RESTRICTED DRUG (CALIFORNIA)- USE ONLY AS DIRECTED.

WARNING: Withdraw from feed 5 days prior to slaughter. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. A milk discard time has not been established for use in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older. Use of more than one product containing neomycin or failure to follow withdrawal times may result in illegal drug residues.

Buyer assumes all responsibility for use, storage and handling of this product. Accelerated Genetics® makes no other claims or warranties expressed or implied.

Accel Calf ADE+ is a registered trademark of and manufactured for Accelerated Genetics, Westby, WI 54667

800-451-9275 | INFO@ACCELGEN.COM | WWW.ACCELGEN.COM



ACCEL CALF ADE+ TYPE B MEDICATED FEED

A calf supplement to be mixed with dry non-medicated milk replacer to make a complete medicated Type C milk replacer ration for calves and beef and non-lactating dairy cattle.

ACTIVE DRUG INGREDIENTS

GUARANTEED ANALYSIS

Crude Protein, Min	20.00%
Crude Fat, Min	20.00%
Crude Fiber, Max	0.15%
Calcium, Min.	0.50%
Calcium, Max	0.75%
Phosphorus, Min.	0.50%
Potassium, Min	1.00%
Salt, Min	1.50%
Salt, Max.	2.00%
Selenium, Min	0.30 ppm
Vitamin A, Min	30,000 IU/lb.
Vitamin D ₂ , Min	10,000 IU/lb.
Vitamin E, Min.	

INGREDIENTS

Dried whey protein concentrate, dried whey, animal fat (preserved with BHA, BHT, citric acid & ethoxyquin), dried skimmed milk, sodium silico aluminate, dicalcium phosphate, calcium carbonate, artificial flavor, iron proteinate, vitamin E supplement, choline chloride, magnesium sulfate, zinc proteinate, selenium yeast, brewer's dried yeast, DL-methionine, manganese proteinate, vitamin A supplement, copper proteinate, vitamin \mathbf{D}_3 supplement, calcium iodate, ascorbic acid, cobalt proteinate, maltodextrin, niacin supplement, calcium pantothenate, vitamin \mathbf{B}_{12} supplement, menadione sodium bisulfite complex (source of vitamin K activity), thiamine mononitrate, riboflavin supplement, folic acid, pyridoxine hydrochloride, biotin.

CAUTION

If symptoms persist after using for 2 or 3 days, consult a veterinarian. Treatment should continue 24 to 48 hours beyond remission of disease symptoms.

FOR USE IN DRY FEEDS ONLY. NOT FOR USE IN LIQUID FEED SUPPLEMENTS. FOR USE IN ANIMAL FEED ONLY. NOT FOR HUMAN USE RESTRICTED DRUG (CALIFORNIA)- USE ONLY AS DIRECTED.

FEEDING AND MIXING DIRECTIONS

For the treatment of bacterial enteritis (scours) caused by *Escherichia coli* and bacterial pneumonia (shipping fever complex) caused by *Pasteurella multocida* susceptible to oxytetracycline; also for the treatment and control of colibacillosis (bacterial enteritis, scours) caused by *Escherichia coli* susceptible to neomycin. Feed to provide each calf *10 mg of oxytetracycline and 10 mg of neomycin sulfate per pound of calf bodyweight daily. Feed continously for 7 to 14 days in place of normal milk replacer.

Mix the entire package (4.34 lbs.) of Accel Calf Ade+ powder into 50 pounds of dry non-medicated calf milk replacer powder to make a Type C complete, medicated milk replacer ration containing 1,000 mg Oxytetracycline and 1,000 mg Neomycin Sulfate per 1.25 pounds of medicated milk replacer powder.

Mix and feed the resulting Type C feed as a complete ration according to the following chart:

DIRECTIONS FOR USE

Calf Weight (lbs.)	*Neomycin & Oxytetracycline (mg/day)	Type C Milk Replacer Powder (ounces)	Water (115-120°F) (quarts)	Feedings Per Day
80	800	8	2	2
100	1000	10	2	2
125	1250	12.5	2.5	2
150	1500	15	3	2

WARNING

Withdraw from feed 5 days prior to slaughter. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. A milk discard time has not been established for use in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older. Use of more than one product containing neomycin or failure to follow withdrawal times may result in illegal drug residues.



Buyer assumes all responsibility for use, storage and handling of this product. Accelerated Genetics® makes no other claims or warranties expressed or implied. Accel Calf Ade+ is manufactured for Accelerated Genetics®, Westby, WI 54667